

Spin Polarized Asymmetric Nuclear Matter and Neutron Star Matter Within the Lowest Order Constrained Variational Method

G.H. Bordbar^{a,b * †} and M. Bigdeli^{a,c}

^a*Department of Physics,*

Shiraz University, Shiraz 71454, Iran[‡],

^b*Research Institute for Astronomy and Astrophysics of Maragha,*

P.O. Box 55134-441, Maragha, Iran

and

^c*Department of Physics,*

Zanjan University, Zanjan, Iran

Abstract

In this paper, we calculate properties of the spin polarized asymmetrical nuclear matter and neutron star matter, using the lowest order constrained variational (LOCV) method with the AV_{18} , $Reid93$, UV_{14} and AV_{14} potentials. According to our results, the spontaneous phase transition to a ferromagnetic state in the asymmetrical nuclear matter as well as neutron star matter do not occur.

* Corresponding author

† E-mail : bordbar@physics.susc.ac.ir

‡ Permanent address

I. INTRODUCTION

The magnetic properties of polarized neutron matter, nuclear matter and neutron star matter have crucial role for studying the possible onset of ferromagnetic transition in the neutron star core. The most interesting and stimulating mechanisms that have been suggested for magnetic source of pulsars, is the possible existence of a phase transition to a ferromagnetic state at densities corresponding to the theoretically stable neutron stars and, therefore, of a ferromagnetic core in the liquid interior of such compact objects. Pulsars are rapidly rotating neutron stars with strong surface magnetic fields in the range of $10^{12} - 10^{13}$ Gauss [1, 2, 3]. Such a possibility has been studied by several authors using different theoretical approaches [4-29], but the results are still contradictory. In most calculations, neutron star matter is approximated by pure neutron matter, as proposed just after the discovery of pulsars. Vidana et al. [21] and Vidana and Bombaci [22] have considered properties of spin polarized neutron matter and polarized asymmetrical nuclear matter using the Brueckner-Hartree-Fock (BHF) approximation by employing three realistic nucleon-nucleon interactions, Nijmegen II, Reid93 and NSC97e respectively. Zuo et al. [25] have also obtained properties of spin polarized neutron and symmetric nuclear matter using same method with AV_{18} potential. The results of those calculations show no indication of ferromagnetic transition at any density for neutron and asymmetrical nuclear matter. Fantoni et al. [20] have calculated spin susceptibility of neutron matter using the Auxiliary Field Diffusion Monte Carlo (AFDMC) method employing the AU6 + UIX three-body potential, and have found that the magnetic susceptibility of neutron matter shows a strong reduction of about a factor 3 with respect to its Fermi gas value. Baldo et al. [26], Akmal et al. [27] and Engvik et al. [28] have considered properties of neutron matter with AV_{18} potential using BHF approximation both for continuous choice (BHFC) and standard choice (BHFG), variational chain summation (VCS) method and lowest order Brueckner (LOB) respectively. On the other hand some calculations, like those of Brownell and Callaway [4] and Rice [5] considered a hard sphere gas model and showed that neutron matter becomes ferromagnetic at $k_F \approx 2.3 fm^{-1}$. Silverstein [8] and Østgaard [9] found that the inclusion of long range attraction significantly increased the ferromagnetic transition density (e.g., Østgaard predicted the transition to occur at $k_F \approx 4.1 fm^{-1}$ using a simple central potential with hard core only for the singlet spin states). Clark [6] and Pearson and Saunier [10]

calculated the magnetic susceptibility for low densities ($k_F \leq 2fm^{-1}$) using more realistic interactions. Pandharipande et al. [11], using the Reid soft-core potential, performed a variational calculation arriving to the conclusion that such a transition was not to be expected for $k_F \leq 5fm^{-1}$. Early calculations of the magnetic susceptibility within the Brueckner theory were performed by Bäckmann and Källman [12] employing the Reid soft-core potential, and results from a correlated basis function calculation were obtained by Jackson et al. [14] with the Reid v6 interaction. A different point of view was followed by Vidaurre et al. [21], who employed neutron-neutron effective interactions of the Skyrme type, finding the ferromagnetic transition at $k_F \approx 1.73 - 1.97fm^{-1}$. Marcos et al. [16] have also studied the spin stability of dense neutron matter within the relativistic Dirac-Hartree-Fock approximation with an effective nucleon-meson Lagrangian, predicting the ferromagnetic transition at several times nuclear matter saturation density. This transition could have important consequences for the evolution of a protoneutron star, in particular for the spin correlations in the medium which do strongly affect the neutrino cross section and the neutrino mean free path inside the star [30].

In our pervious works, the properties of unpolarized asymmetrical nuclear matter have been considered by us using the lowest order constrained variational (LOCV) method [31] with the AV_{18} potential [32]. We have found that the energy per particle of nuclear matter alter monotonically and linearly by quadratic asymmetrical parameter.

Recently, we have computed the properties of polarized neutron matter[33] and polarized symmetrical nuclear matter[34] such as total energy, magnetic susceptibility, pressure, etc using the microscopic calculations employing LOCV method with the AV_{18} potential. We have also concluded that the spontaneous phase transition to a ferromagnetic state in the neutron and asymmetrical nuclear matter does not occur. In this work, we intend to calculate the properties of spin polarized asymmetrical nuclear matter and neutron star matter using the LOCV method employing the AV_{18} [32], *Reid93* [35], *UV₁₄* [36] and AV_{14} [37] potentials.

II. LOCV METHOD

The LOCV method which was developed several years ago is a useful tool for the determination of the properties of neutron, nuclear and asymmetric nuclear matter at zero and finite temperature. The LOCV method is a fully self-consistent formalism and it does not

bring any free parameters into calculation. It employs a normalization constraint to keep the higher order term as small as possible. The functional minimization procedure represents an enormous computational simplification over unconstrained methods that attempt to go beyond lowest order [31, 38, 39].

In this method we consider a trial many-body wave function of the form

$$\psi = F\phi, \quad (1)$$

where ϕ is the uncorrelated ground state wave function (simply the Slater determinant of plane waves) of A independent nucleon and $F = F(1 \cdots A)$ is an appropriate A -body correlation operator which can be replaced by a Jastrow form i.e.,

$$F = S \prod_{i>j} f(ij), \quad (2)$$

in which S is a symmetrizing operator. We consider a cluster expansion of the energy functional up to the two-body term,

$$E([f]) = \frac{1}{A} \frac{\langle \psi | H \psi \rangle}{\langle \psi | \psi \rangle} = E_1 + E_2. \quad (3)$$

The one-body term E_1 is total kinetic energy of the system. The two-body energy E_2 is

$$E_2 = \frac{1}{2A} \sum_{ij} \langle ij | \nu(12) | ij - ji \rangle, \quad (4)$$

where

$\nu(12) = -\frac{\hbar^2}{2m} [f(12), [\nabla_{12}^2, f(12)]] + f(12)V(12)f(12)$, $f(12)$ and $V(12)$ are the two-body correlation and potential. For the two-body correlation function, $f(12)$, we consider the following form [31, 38]:

$$f(12) = \sum_{k=1}^3 f^{(k)}(12) O^{(k)}(12), \quad (5)$$

where, the operators $O^{(k)}(12)$ are given by

$$O^{(k=1-3)}(12) = 1, \left(\frac{2}{3} + \frac{1}{6}S_{12}\right), \left(\frac{1}{3} - \frac{1}{6}S_{12}\right), \quad (6)$$

and S_{12} is the tensor operator.

$$S_{12} = 3(\sigma_1 \cdot \hat{\mathbf{r}})(\sigma_2 \cdot \hat{\mathbf{r}}) - \sigma_1 \cdot \sigma_2 \quad (7)$$

Now, we can minimize the two-body energy Eq.(4), with respect to the variations in the function $f_\alpha^{(i)}$ but subject to the normalization constraint [31],

$$\frac{1}{A} \sum_{ij} \langle ij | h_{S_z}^2 - f^2(12) | ij \rangle_a = 0, \quad (8)$$

where in the case of spin polarized neutron matter the function $h_{S_z}(r)$ is defined as

$$\begin{aligned} h_{S_z}(r) &= \left[1 - \frac{9}{\nu} \left(\frac{J_J^2(k_F^i)}{k_F^i} \right)^2 \right]^{-1/2} ; S_z = \pm 1 \\ &= 1 ; S_z = 0 \end{aligned} \quad (9)$$

here ν is the degeneracy of the system. From the minimization of the two-body cluster energy, we get a set of coupled and uncoupled differential equations the same as presented in Ref. [31]. We can get correlation functions by solving the differential equations and in turn these functions lead to the two-body energy.

III. SPIN POLARIZED ASYMMETRICAL NUCLEAR MATTER

Spin polarized asymmetrical nuclear matter is an infinite system that is composed of spin up and spin down neutrons with densities $\rho_n^{(1)}$ and $\rho_n^{(2)}$ respectively, and spin up and spin down protons with densities $\rho_p^{(1)}$ and $\rho_p^{(2)}$. The total densities for neutrons (ρ_n), protons (ρ_p), and nucleons (ρ) are given by:

$$\rho_p = \rho_p^{(1)} + \rho_p^{(2)}, \quad \rho_n = \rho_n^{(1)} + \rho_n^{(2)}, \quad \rho = \rho_p + \rho_n \quad (10)$$

Labels 1 and 2 are used instead of spin up and spin down nucleons, respectively. One can use the following parameter to identify a given spin polarized state of the system,

$$\delta_p = \frac{\rho_p^{(1)} - \rho_p^{(2)}}{\rho}, \quad \delta_n = \frac{\rho_n^{(1)} - \rho_n^{(2)}}{\rho} \quad (11)$$

δ_p and δ_n are proton and neutron spin asymmetry parameters, respectively. Total polarization defined as,

$$\delta = \delta_n + \delta_p \quad (12)$$

For the unpolarized case, we have $\delta_n = \delta_p = 0$.

Asymmetry parameter describes isospin asymmetry of the system and is defined as,

$$\beta = \frac{\rho_n - \rho_p}{\rho}. \quad (13)$$

Pure neutron matter is totally asymmetrical nuclear matter with $\beta = 1$ and symmetrical nuclear matter has $\beta = 0$. In the case of spin polarized asymmetrical nuclear matter, the energy per particle given by

$$E(\rho, \delta, \beta) = E_1(\rho, \delta, \beta) + E_2(\rho, \delta, \beta). \quad (14)$$

where E_1 , the kinetic energy contribution given by,

$$E_1(\rho, \delta, \beta) = \frac{3}{20} \frac{\hbar^2 k_F^2}{2m} \left\{ (1 + \beta + 2\delta_n)^{\frac{5}{3}} + (1 + \beta - 2\delta_n)^{\frac{5}{3}} \right. \\ \left. + (1 - \beta + 2\delta_p)^{\frac{5}{3}} + (1 - \beta - 2\delta_p)^{\frac{5}{3}} \right\}, \quad (15)$$

where $k_F = (3/2\pi^2\rho)^{1/3}$ is Fermi momentum of unpolarized symmetrical nuclear matter. In expression (14), the two body energy can be calculated by the semi-empirical mass formula [31],

$$E_2(\rho, \delta, \beta) = \beta^2 E_{2neum} + (1 - \beta^2) E_{2snucm}, \quad (16)$$

where the powers higher than quadratic are neglected. E_{2neum} is polarized pure neutron matter potential energy, and E_{2snucm} is polarized symmetrical nuclear matter potential energy that both of them have been determined using the LOCV method employing a microscopic point of view [33, 34].

The energy per particle of the polarized asymmetrical nuclear matter versus density at different values of the spin polarization and isospin asymmetry parameter have been shown in Figs. 1-4 for the AV_{18} , $Reid93$, UV_{14} and AV_{14} potentials, respectively. As it can be seen from these figures, the energy of polarized asymmetrical nuclear matter for various values of isospin asymmetry parameters become more repulsive by increasing the polarization for all relevant densities. There is no crossing of the energy curves of different polarizations. Reversibly, by increasing density, the difference between the energy of nuclear matter at different polarization becomes more sizable. Indeed, the ground state of the system is that of unpolarized matter in all ranges of density and isospin asymmetry considered. This behavior is common for all potentials used in our calculations.

Another quantities that help us to understand ferromagnetic phase transition, are magnetic susceptibility and Landau parameter. In what follows, we want to consider these two quantities. The magnetic susceptibility, χ , which characterizes the response of a system to the magnetic field and gives a measure of the energy required to produce a net spin alignment in the direction of the magnetic field, is defined by

$$\chi = \left(\frac{\partial M}{\partial H} \right)_{H=0}, \quad (17)$$

where M is the magnetization of the system per unit volume and H is the magnetic field. We have calculated the magnetic susceptibility of the polarized asymmetrical nuclear matter in the ratio χ/χ_F form. By using the Eq. (17) and some simplification, the ratio of χ to the magnetic susceptibility for a degenerate free Fermi gas χ_F can be written as

$$\frac{\chi}{\chi_F} = \frac{2}{3} \frac{E_F}{\left(\frac{\partial^2 E}{\partial \delta^2} \right)_{\delta=0}}, \quad (18)$$

where $E_F = \hbar^2 k_F^2 / 2m$ is the Fermi energy. After doing some algebra the ratio χ/χ_F takes the form,

$$\frac{\chi}{\chi_F} = \left[1 + \frac{2 \left(\beta^2 \left\{ 2^{\frac{2}{3}} (\chi_F/\chi)_{neum} - (\chi_F/\chi)_{snucm} \right\} + (\chi_F/\chi)_{snucm} + \beta^2 (1 - 2^{\frac{2}{3}}) - 1 \right)}{(1 + \beta)^{5/3} + (1 - \beta)^{5/3}} \right]^{-1} \quad (19)$$

where $(\chi/\chi_F)_{neum}$ and $(\chi/\chi_F)_{snucm}$ is magnetic susceptibility of pure neutron matter [33], and asymmetry nuclear matter [34], respectively. In Fig. 5, we have plotted the ratio χ/χ_F as a function of density at several values of the isospin asymmetry for the AV_{18} , $Reid93$, UV_{14} and AV_{14} potentials. For all used potentials, we see that the value of the ratio χ/χ_F decreases monotonically with the density, even at high densities. It shows, there is no magnetic instability for any values of asymmetry parameter. If such an instability existed, the value of ratio χ/χ_F changes suddenly unlike previous treatment.

As it is known, the Landau parameter, G_0 , describes the spin density fluctuation in the effective interaction. G_0 is simply related to the magnetic susceptibility by the relation

$$\frac{\chi}{\chi_F} = \frac{m^*}{1 + G_0}, \quad (20)$$

where m^* is the effective mass. A magnetic instability would require $G_0 < -1$. Our results for the Landau parameter have been presented in the Fig. 6 for the AV_{18} , $Reid93$, UV_{14} and AV_{14} potentials. It is seen that the value of G_0 is always positive and monotonically

increases by increasing the density. This shows that for all used potentials, the spontaneous phase transition to a ferromagnetic state in the asymmetrical nuclear matter does not occur.

The equation of state of polarized asymmetrical nuclear matter, $P(\rho, \beta, \delta)$, can be obtained using

$$P(\rho, \beta, \delta) = \rho^2 \frac{\partial E(\rho, \beta, \delta)}{\partial \rho}. \quad (21)$$

In Fig. 7, we have shown the pressure of asymmetrical nuclear matter as a function of density (ρ) at different polarizations for various choice of asymmetry parameter (β) with the AV_{18} potential. This figure shows that the equation of state becomes stiffer by increasing the polarization for all isospin asymmetry.

IV. POLARIZED NEUTRON STAR MATTER

Indeed the neutron star matter is charge neutral infinite system that is mixture of asymmetrical nuclear matter and leptons, specially electrons and muons. The energy per particle of polarized neutron star matter, E_{nsm} can be written as,

$$E_{nsm} = E + E_l. \quad (22)$$

where E is the nucleonic energy contribution which given by Eq. (14) and E_l is leptonic energy contribution obtained as follows,

$$E_l = \sum_{i=e, \mu} \sum_{k \leq k^{(F)}} [(m_i c^2)^2 + \hbar^2 c^2 k^2]^{1/2}. \quad (23)$$

After some simplification the above equation leads to

$$E_l = \frac{3}{8} \sum_{i=e, \mu} \frac{\rho_i}{\rho} \frac{m_i c^2}{x_i^{(F)^3}} \left[x_i^{(F)} (2x_i^{(F)^2} + 1) \sqrt{x_i^{(F)^2} + 1} - \sinh^{-1} x_i^{(F)} \right], \quad (24)$$

where $x_i^{(F)} = \hbar k_i^{(F)} / m_i c$. The conditions of charge neutrality and beta equilibrium impose the following constraints on the calculation of energy of neutron star matter [1],

$$\mu_n = \mu_p + \mu_e \quad \mu_e = \mu_\mu. \quad (25)$$

$$\rho_p = \rho_e + \rho_\mu. \quad (26)$$

In Fig. 8, we have shown the energy per particle of the neutron star matter at various values of spin polarization as a function of density for the AV_{18} , $Reid93$, UV_{14} and AV_{14} potentials. The same as Figs. 1-4, this figure have also shown no phase transition to a ferromagnetic state.

The treatment of magnetic susceptibility and landau parameter of neutron star matter versus density for different polarization, have been shown in Figs. 9 and 10, respectively. This treatments are the same as the treatments of the polarized neutron matter, symmetrical and asymmetrical nuclear matter, and it does not show any magnetic instability for the neutron star matter. This behavior has been observed for all potentials used in our calculations.

For the AV_{18} potential, the pressure of neutron star matter have been presented as a function of density ρ at different polarizations in Fig. 11. We see that the equation of state of neutron star matter becomes stiffer by increasing the polarization.

V. SUMMARY AND CONCLUSIONS

The purpose of this paper was to calculate the properties of the spin polarized asymmetrical nuclear matter and neutron star matter employing the lowest order constrained variational technique with the AV_{18} , $Reid93$, UV_{14} and AV_{14} nucleon-nucleon potentials. After introducing the LOCV method briefly, we calculated the energy per particle of the polarized asymmetrical nuclear matter and showed that the force becomes more repulsive by increasing the polarization for all relevant densities. No crossing of the energy curves was observed and the ground state of the system was found to be that of the unpolarized matter in all ranges of densities and isospin asymmetry. The magnetic susceptibility was computed and shown to remain almost constant by increasing the asymmetry parameter. For the polarized neutron star matter, we showed that there is no magnetic instability and the equation of state becomes stiffer by increasing the polarization.

Acknowledgments

This work has been supported financially by Research Institute for Astronomy and Astrophysics of Maragha. One os us (G.H. Bordbar) wish to thanks Shiraz University Research

Council.

-
- [1] S. Shapiro and S. Teukolsky, *Black Holes, White Dwarfs and Neutron Stars*, (Wiley-New York, 1983).
 - [2] F. Pacini, *Nature* (London) **216** (1967) 567.
 - [3] T. Gold, *Nature* (London) **218** (1968) 731.
 - [4] D. H. Brownell and J. Callaway, *Nuovo Cimento* **B 60** (1969) 169.
 - [5] M. J. Rice, *Phys. Lett.* **A 29** (1969) 637.
 - [6] J. W. Clark and N. C. Chao, *Lettere Nuovo Cimento* **2** (1969) 185.
 - [7] J. W. Clark, *Phys. Rev. Lett.* **23** (1969) 1463.
 - [8] S. D. Silverstein, *Phys. Rev. Lett.* **23** (1969) 139.
 - [9] E. Østgaard, *Nucl. Phys.* **A 154** (1970) 202.
 - [10] J. M. Pearson and G. Saunier, *Phys. Rev. Lett.* **24** (1970) 325.
 - [11] V. R. Pandharipande, V. K. Garde and J. K. Srivastava, *Phys. Lett.* **B 38** (1972) 485.
 - [12] S. O. Backman and C. G. Kallman, *Phys. Lett.* **B 43** (1973) 263.
 - [13] P. Haensel, *Phys. Rev.* **C 11** (1975) 1822.
 - [14] A. D. Jackson, E. Krotscheck, D. E. Meltzer and R. A. Smith, *Nucl. Phys.* **A 386** (1982) 125.
 - [15] M. Kutschera and W. Wojcik, *Phys. Lett.* **B 223** (1989) 11.
 - [16] S. Marcos, R. Niembro, M. L. Quelle and J. Navarro, *Phys. Lett.* **B 271** (1991) 277.
 - [17] P. Bernardos, S. Marcos, R. Niembro, M. L. Quelle, *Phys. Lett.* **B 356** (1995) 175.
 - [18] A. Vidaurre, J. Navarro and J. Bernabeu, *Astron. Astrophys.* **135** (1984) 361.
 - [19] M. Kutschera and W. Wojcik, *Phys. Lett.* **B 325** (1994) 271.
 - [20] S. Fantoni, A. Sarsa and K. E. Schmidt, *Phys. Rev. Lett.* **87** (2001) 181101.
 - [21] I. Vidana, A. Polls and A. Ramos, *Phys. Rev.* **C 65** (2002) 035804.
 - [22] I. Vidana and I. Bombaci, *Phys. Rev.* **C 66** (2002) 045801.
 - [23] W. Zuo, U. Lombardo and C.W. Shen, in *Quark-Gluon Plasma and Heavy Ion Collisions*, Ed. W.M. Alberico, M. Nardi and M.P. Lombardo, World Scientific, p. 192 (2002).
 - [24] A. A. Isayev and J. Yang, *Phys. Rev.* **C 69** (2004) 025801.
 - [25] W. Zuo, U. Lombardo and C. W. Shen, nucl-th/0204056.
W. Zuo, C. W. Shen and U. Lombardo, *Phys. Rev* **C 67** (2003) 037301.

- [26] M. Baldo, G. Giansiracusa, U. Lombardo and H. Q. Song, *Phys. Lett.* **B 473** (2000) 1.
- [27] A. Akmal, V. R. Pandharipande and D. G. Ravenhall, *Phys. Rev.* **C 58** (1998) 1804.
- [28] L. Engvik et al., *Nucl. Phys.* **A 627** (1997) 85.
- [29] A.Rios, A. Polls and I. Vidana, *Phys. Rev.* **C 71**(2005) 055802.
- [30] J. Navarro, E. S. Hernandez and D. Vautherin, *Phys. Rev.* **C 60** (1999) 045801.
- [31] G. H. Bordbar and M. Modarres, *Phys. Rev.* **C 57** (1998) 714.
- [32] R. B. Wiringa, V. Stoks and R. Schiavilla, *Phys. Rev.* **C 75** (1995) 38.
- [33] G. H. Bordbar and M. Bigdeli, *Phys. Rev.* **C 75** (2007) 045804.
- [34] G. H. Bordbar and M. Bigdeli, *Phys. Rev.* **C 76** (2007) 035803.
- [35] V.G.J. Stoks, R.A.M. Klomp, C.P.F. Terheggen and J.J. de Swart, *Phys. Rev.* **C 49** (1994) 2950.
- [36] I.E. Lagaris and V.R. Pandharipande, *Nucl. Phys.* **A 359** (1981) 331.
- [37] R.B. Wiringa, R.A. Smith and T.L. Ainsworth, *Phys. Rev.* **C 29** (1984) 1207.
- [38] G. H. Bordbar, M. Modarres, *J. Phys. G: Nucl. Part. Phys.* **23** (1997) 1631.
- [39] J. C. Owen, R. F. Bishop, and J. M. Irvine, *Nucl. Phys.* **A 277** (1977) 45.




FIG. 1: The energy per particle of the polarized asymmetrical nuclear matter versus density(ρ) for different values of the spin polarization (δ) with the AV_{18} potential at $\beta = 0.0$ (a), $\beta = 0.3$ (b), $\beta = 0.6$ (c), $\beta = 1.0$ (d).




FIG. 2: The energy per particle of the polarized asymmetrical nuclear matter versus density(ρ) for different values of the spin polarization (δ) with the Reid93 potential at $\beta = 0.0$ (a), $\beta = 0.3$ (b), $\beta = 0.6$ (c), $\beta = 1.0$ (d).




FIG. 3: The energy per particle of the polarized asymmetrical nuclear matter versus density(ρ) for different values of the spin polarization (δ) with the UV_{14} potential at $\beta = 0.0$ (a), $\beta = 0.3$ (b), $\beta = 0.6$ (c), $\beta = 1.0$ (d).




FIG. 4: The energy per particle of the polarized asymmetrical nuclear matter versus density(ρ) for different values of the spin polarization (δ) with the AV_{14} potential at $\beta = 0.0$ (a), $\beta = 0.3$ (b), $\beta = 0.6$ (c), $\beta = 1.0$ (d).




FIG. 5: The magnetic susceptibility of the polarized asymmetrical nuclear matter as the function of density (ρ) for different values of asymmetry (β) with the AV_{18} (a), Reid93 (b), UV_{14} (c) and AV_{14} (d) potentials.




FIG. 6: The Landau parameter of the polarized asymmetrical nuclear matter as the function of density (ρ) for different values of asymmetry (β) with the AV_{18} (a), $Reid93$ (b), UV_{14} (c) and AV_{14} (d) potentials.




FIG. 7: The equation of state of polarized asymmetrical nuclear matter for different values of the spin polarization (δ) with the AV_{18} potential at $\beta = 0.0$ (a), $\beta = 0.3$ (b), $\beta = 0.6$ (c), $\beta = 1.0$ (d).




FIG. 8: The energy per particle of the polarized neutron star matter versus density (ρ) for different values of the spin polarization (δ) with the AV_{18} (a), Reid93 (b), UV_{14} (c) and AV_{14} (d) potentials.




FIG. 9: The magnetic susceptibility of the polarized neutron star matter versus density (ρ) with the AV_{18} , Reid93, UV_{14} and AV_{14} potentials.




FIG. 10: The Landau parameter, G_0 , of polarized neutron star matter as function of density(ρ) with the AV_{18} , Reid93, UV_{14} and AV_{14} potentials.




FIG. 11: The equation of state of polarized neutron star matter for different values of the spin polarization (δ) with the AV_{18} potential .